arXiv:1608.04646v1 [hep-ph] 16 Aug 2016

Saturation effects in low-x DIS structure functions and related hadronic total cross sections

Francesco Giovanni Celiberto[1,2,⋆], *Laszlo* Jenkovszky[3,⋆⋆], and *Volodymyr* Myronenko[4,⋆⋆⋆]

[1] *Dipartimento di Fisica, Università della Calabria, Arcavacata di Rende, 87036 Cosenza, Italy*
[2] *Istituto Nazionale di Fisica Nucleare, Gruppo Collegato di Cosenza, Arcavacata di Rende, 87036 Cosenza, Italy*
[3] *Bogolyubov Institute for Theoretical Physics (BITP), Ukrainian National Academy of Sciences 14-b, Metrologicheskaya str., Kiev, 03680, Ukraine*
[4] *Deutsches Elektronen-Synchrotron DESY, Hamburg, Germany*

Abstract.
High-energy nucleon total cross sections are related to low-x DIS structure functions by using the additive quark model.

In the additive quark model, the hadron-hadron total cross section can be written as a product of the cross sections of the constituents, σ_{qq} [1, 2], *e.g.*

$$\sigma(s)^t_{pp} = \sigma_{qq}[n_V + n_S(s)]^2, \tag{1}$$

where n_V is the number of valence quarks and $n_S(s)$ is that of sea quarks, their number increasing with energy.

It was suggested in Refs. [1, 2] that the increasing number of sea quarks is related to the Bjorken scaling-violating contribution to the deep inelastic lepton-hadron structure function (DIS SF), namely to the momentum fraction of the relevant quarks given by the integral over the DIS structure function $F_2(x, Q^2)$. In Ref. [1] a simple model for the DIS structure function, known at those times, was used, resulting in the following expression for the total cross section, compatible with the data

$$\sigma(s)^t_{pp} = \sigma_{qq} n_V^2 (1 + 0.0.16\, \ell n(s/Q_0^2)), \tag{2}$$

where σ_{qq} is a free parameter, Q_0^2 was fitted to the DIS data, and $n_V = 3$.

In Ref. [2] the DIS SF was related to hadronic cross sections by means of finite-energy sum rules in Q^2.

The number of quarks in a reaction can be calculated from the SF by means of sum rules. see e.g. [3, 4].

In Ref. [5] following ansatz for the small-x singlet part (labelled by the upper index $S, 0$) of the proton structure function, interpolating between the soft (VMD, Pomeron) and hard (GLAP evolution) regimes was proposed:

⋆ e-mail: francescogiovanni.celiberto@fis.unical.it
⋆⋆ e-mail: jenk@bitp.kiev.ua
⋆⋆⋆ e-mail: volodymyr.myronenko@desy.de



Figure 1. Slope of the structure function $F_2(x, Q^2)$.

$$F_2^{(S,0)}(x, Q^2) = A\left(\frac{Q^2}{Q^2 + a}\right)^{1+\widetilde{\Delta}(Q^2)} e^{\Delta(x,Q^2)}, \tag{3.1}$$

with the "effective power"

$$\widetilde{\Delta}(Q^2) = \epsilon + \gamma_1 \ell n\left(1 + \gamma_2 \ell n\left[1 + \frac{Q^2}{Q_0^2}\right]\right), \tag{3.2}$$

and

$$\Delta(x, Q^2) = \left(\widetilde{\Delta}(Q^2)\ell n\frac{x_0}{x}\right)^{f(Q^2)}, \tag{3.3}$$

where

$$f(Q^2) = \frac{1}{2}\left(1 + e^{-Q^2/Q_1^2}\right). \tag{3.4}$$

At small and moderate values of Q^2, the exponent $\widetilde{\Delta}(Q^2)$ (3.2) may be interpreted as a Q^2-dependent "effective Pomeron intercept", as shown in Fig. 1.

The function $f(Q^2)$ has been introduced in order to provide for the transition from the Regge behaviour, where $f(Q^2) = 1$, to the asymptotic solution of the GLAP evolution equation, where $f(Q^2) = 1/2$.

In Ref. [5] the above singlet SF was appended by a non-singlet part, important at large values of x. The parameters were fitted to the DIS data in a wide range of x and Q^2.The values of the fitted parameters are: $A = 0.1623$, $a = 0.2916$ GeV^2, $\gamma_2 = 0.01936$, $Q_0^2 = 0.1887$ GeV^2, $Q_1^2 = 916.1$ GeV^2; $x_0 = 1$, $\epsilon = 0.08$, $\gamma_1 + 2.4$ were fixed (by QCD-related arguments). The resulting fits and more details can be found in Ref. [5].

The proton-proton total cross section is cast by integrating Eqs. (3) between $x = 0$ and $x = 1$. At high energies, only the singlet part of the SF, Eqs. (3) (the "Pomeron") is relevant. Integration can be



Figure 2. HHT (top) and HERAPDF2.0 (bottom) NLO predictions of the F_2 structure function at $Q^2_{\text{min}} = 3.5\,\text{GeV}^2$, compared to extracted values. For more details, see Ref. [7].

performed numerically. The result is in reasonable agreement with the data on pp total cross sections, including those from the LHC.

The operator-product expansion beyond leading twist has diagrams in which two, three or four gluons may be exchanged in the t-channel such that these gluons may be viewed as recombining. This recombination could lead to gluon saturation. The colour-dipole framework also inspired a phenomenological model of saturation by Golec–Biernat–Wüsthoff (GBW), in which the onset of saturation is characterised as the transition from a "soft" to a "hard" scattering regime. This occurs along a "critical line" in the x_{Bj}, Q^2 plane.

Recently new results on low-x DIS parton distrubutions (PDF and SF) have appeared [6, 7]; they show an intriguing change of regime towards smallest values of x (Fig. 2) - possible saturation effect?

We intend to investigate its impact on the asymptotic behaviour of the total cross sections.

Acknowledgement

L.J. thanks the Organizers of the Quarks2016 conference for their warm hospitality and support.

References

[1] L.L. Jenkovszky and B.V. Struminsky: *Violation of scale invariance in deep-inelastic lepton-hadron scattering and rise of hadronic cross-sections*, ITP-77-37E Preprint, Kiev, 1977.

[2] L.L. Jenkovszky and B.V. Struminsky: Yadernaya Fizika (English translation: Sov. J. Nucl. Phys.), **38**(1983), 1568; see also: http://inspirehep.net/record/1456006/files/$mpi_a t_l hc_2$015.pdf; http://indico.ictp.it/event/a14280/ , p.171.

[3] F.E. Close, *An introduction to quarks and partons*, Academic Press Inc. (London) Ltd., 1979.

[4] R.G. Roberts, *The structure of the proton*, Cambridge University Press, 1990.

[5] P. Desgrolard, Laszlo L. Jenkovszky, F. Paccanoni, *Interpolating between soft and hard dynamics in deep inelastic scattering*. LYCEN-9808, Eur. Phys. J. C7 (1999) 263-270; hep-ph/9803286.

[6] H. Abramowicz *et al.* [H1 and ZEUS Collaborations], *Combination of measurements of inclusive deep inelastic $e^{\pm}p$ scattering cross sections and QCD analysis of HERA data*, Eur. Phys. J. C **75** (2015) no.12, 580 doi:10.1140/epjc/s10052-015-3710-4 [arXiv:1506.06042 [hep-ex]].

[7] I. Abt, A. M. Cooper-Sarkar, B. Foster, V. Myronenko, K. Wichmann and M. Wing, *Study of HERA ep Data at Low Q^2 and Low x_{B_j} and the Need for Higher-Twist Corrections to Standard pQCD Evolution*, arXiv:1604.02299 [hep-ph].